



Eccentricity of selected eclipsing systems

W.Ogłóza, J.M.Kreiner, G.Stachowski, B.Zakrzewski
Mt. Suhora Observatory, Cracow Pedagogical University, Poland

Z.Mikulášek, M.Zejda
Masaryk University, Brno, Czech Republic



The study of orbital eccentricity and apsidal motion are very important for verification of theories of evolution of binary systems and of the internal structure of stars. One of the methods used to study long time effects is (O-C) analysis of times of minima.

This poster presents preliminary results of the selection of GCVS eclipsing systems with phase-shifted secondary minima. At present, the Cracow database contains 19493 individual minima of 5931 stars. Examination of the (O-C) diagrams shows about 256 stars with a significant phase shift of the secondary minimum, interpreted as the effect of an eccentric orbit. The poster shows 98 selected systems (with 5800 times of minima) for which the primary and secondary minima were observed at least 4 times over well separated epochs (e.g. the diagram for V799 Cas). About (56%) of those systems show the effect of rotation of the line of apsides (see the diagram for the well known system Y Cyg).



The orbital period of the eccentric orbit binaries is in the range of ~ 1.2 days to ~ 20 days (except OW Gem $P \approx 1200d$, V883 Mon $P \approx 50d$, VW Peg $P \approx 21d$, and LV Her $P \approx 18d$), but for long-period systems the number of minima is small.

There are a few short-period variables which show an apparent phase shift of the secondary minimum (LV Vir $P=0.4094$, TY Men $P=0.4617$, V407 Peg $P=0.6369$, LR Car $P=0.8529$, GI Boo $P=1.0335$, DU Boo $P=1.0559$), however this could be the effect of an asymmetrically-shaped minimum.

For several systems (V947 Cyg, RW Lac, RU Mon, Zeta Phe, V1094 Tau, AO Vel, DR Vul) the effect of an eccentric orbit on the (O-C) diagram is present together with other phenomena, such as constant period change or the influence of a third body.

The full list of eccentric systems and up-to-date linear elements for primary and secondary minima will be continuously published at:

www.as.up.krakow.pl/ephem



The XY axes for all diagrams represent the same parameters. Types of observations are represented by the following symbols for primary and secondary minima:

- CCD: ■ □
- Photoelectric: ● ○
- Visual: • ◦
- Photographic: ■ □
- Skypatrol: + x

