



3rd International Olympiad on Astronomy and Astrophysics

Theoretical Competition

Long Problems

Problem 16: High Altitude Projectile (45 points)

A projectile which starts from the surface of the Earth at the sea level is launched with the initial speed of $v_0 = \sqrt{GM/R}$ and with the projecting angle (with respect to the local horizon) of $\theta = \frac{\pi}{6}$. M and R are the mass and radius of the Earth respectively. Ignore the air resistance and rotation of the Earth.

- Show that the orbit of the projectile is an ellipse with a semi-major axis of $a = R$.
- Calculate the highest altitude of the projectile with respect to the Earth surface (in the unit of the Earth radius).
- What is the range of the projectile (distance between launching point and falling point) in the units of the earth radii?
- What is eccentricity (e) of this elliptical orbit?
- Find the time of flight for the projectile.

Problem 17: Apparent number density of stars in the Galaxy (45 points)

Let us model the number density of stars in the disk of Milky Way Galaxy with a simple exponential function of $n = n_0 \exp\left(-\frac{r-R_0}{R_d}\right)$, where r represents the distance from the center of the Galaxy, R_0 is the distance of the Sun from the center of the Galaxy, R_d is the typical size of disk and n_0 is the stellar density of disk at the position of the Sun. An astronomer observes the center of the Galaxy within a small field of view. We take a particular type of Red giant stars (red clump) as the standard candles for the observation with approximately constant absolute magnitude of $M = -0.2$,

- Considering a limiting magnitude of $m = 18$ for a telescope, calculate the maximum distance that telescope can detect the red clump stars. For simplicity we ignore the presence of interstellar medium so there is no extinction.
- Assume an extinction of 0.7 mag/kpc for the interstellar medium. Repeat the calculation as done in the part (a) and obtain a rough number for the maximum distance these red giant stars can be observed.
- Give an expression for the number of these red giant stars per magnitude within a solid angle of Ω that we can observe with apparent magnitude in the range of m and $m + \Delta m$, (i.e. $\frac{\Delta N}{\Delta m}$). Red giant stars contribute fraction f of overall stars. In this part assume no extinction in the interstellar medium as part (a). Assume the size of the disk is larger than maximum observable distance.